

# Introduction to Gear4 Physics Component

Training course at International User Conference  
on Medicine and Biology applications  
Bordeaux, 8-11 October 2013  
V. Ivanchenko

# Outline

---

- General interface to Geant4 physics
  - Adaptation of **Marc Verderi** original lecture
- Geant4 cuts
- Cuts per G4Region


# Geant4 interface to physics

---

- The `G4ParticleDefinition` interface
- The `G4VProcess` class process interface
- The `G4ProcessManager` class


# Geant4 tracking

- **G4Track** is the object “pushed” step by step by the tracking :



- Moving by one step is the responsibility of the “stepping”
  - Which is the core engine of the “tracking” machinery
- These moves/steps have to be physically meaningful
  - And the stepping invokes physics to realize them
- **This physics is attached to the G4Track, let's see how.**

# From G4Track to processes



# G4VProcess: 3 kind of actions

- Abstract class defining the common interface of **all processes** in Geant4:

- Used by all processes
  - including transportation, etc...
- Defined in `source/processes/management`

- **Three kinds of actions:**

- **AtRest** actions:


- Decay,  $e^+$  annihilation ...

- **AlongStep** actions:

- To describe continuous (inter)actions, occurring along the path of the particle, like ionisation;

- **PostStep** actions:

- For describing point-like (inter)actions, like decay in flight



# G4VProcess : actions summary

---

- The virtual « **action** » methods are following:
  - **AtRestGetPhysicalInteractionLength()** ,  
**AtRestDoIt()** ;
  - **AlongStepGetPhysicalInteractionLength()** ,  
**AlongStepDoIt()** ;
  - **PostStepGetPhysicalInteractionLength()** ,  
**PostStepDoIt()** ;
- Other important virtual method:
  - **G4bool IsApplicable(const G4ParticleDefinition &);**
    - Used to check if a process can handle the given particle type
    - It is called by the kernel when you set up your physics list

# G4VProcess: extensions

---

- A process can implement **any combination** of the three `AtRest`, `AlongStep` and `PostStep` actions:
  - `decay = AtRest + PostStep`
- **If you plan to implement your own process:**
  - A set of intermediate classes exist implementing various combinations of actions:
    - For example:
      - `G4VDiscreteProcess`: only `PostStep` actions
      - `G4VContinuousDiscreteProcess`: `AlongStep + PostStep` actions



# G4ProcessManager

- **G4ProcessManager** maintains three vectors of actions :
  - One for the `AtRest` methods of the particle;
  - One for the `AlongStep` ones;
  - And one for the `PostStep` actions.
- .
- **Note that the ordering of processes provided by/to the G4ProcessManager vectors is relevant and used by the stepping**
  - There are few critical points you should be aware of
    - **Multiple scattering can shift end point of a step and step length**
    - **Scintillation, Cerenkov and some other processes assuming that step and energy deposition at the step are defined**

# Adding a process in physics list

---

- Get the process manager of the particle:

```
G4PhysicsListHelper* helper =  
    G4PhysicsListHelper::GetPhysicsListHelper();  
G4ParticleDefinition* electron = G4Electron::Electron();
```


- Add the process:

```
helper->RegisterProcess(new G4eIonisation, electron);
```

- There is well defined order of processes
- G4PhysicsListhelper is responsible for the correct ordering

# About process ordering

- **The most strong rule for multiple-scattering and transportation.**
- In your physics list, you should **always** have, for the ordering of the **AlongGetPhysicalInteractionLength (...)** methods:
  - Transportation last
    - For all particles
  - Multiple scattering second last
    - For charged particles only
      - assuming **n** processes  
[n-2] ...  
[n-1] multiple scattering  
[n] transportation
- Why ?
  - Processes return a « true path length »;
  - The multiple scattering folds up this length into a **shorter** « geometrical » path length;
  - Based on this new length, the transportation can geometrically limits the step.



# Displaying processes and particles

- When your application has started and when the run manager has been initialized, you can:
  - Check the physics processes attached and their ordering:
    - `/particle/select e-`
    - `/particle/processes/dump`
  - Check what particles exist:
    - `/particle/list`
  - Check a particle property:
    - `/particle/select e-`
    - `/particle/property/dump`
  - Please type “help” to get the full set of commands

---

# Geant4 cuts

# Geant4 approach for cuts


---

- All particles are tracked until it is killed by one of Geant4 process, for example:
  - Out of world volume
  - Inelastic interaction
  - Decay
- If kinetic energy is zero and there is no processes AtRest the particle is killed by stepping manager
- Geant4 by default has no tracking cut but only unique cut in range
  - Physically this means required spatial accuracy of simulation
  - This is the main difference between Geant4 and other simulation tools


# Bremsstrahlung

- **Bremsstrahlung** spectrum grows to low energy as  $1/k$ 
  - $k$  is the gamma energy
- **Low energy gammas** have very small absorption length
- Simulation of all low-energy gammas is **non-effective**
- **Cuts/production threshold** are used in all Monte Carlo codes
- **Gamma emission** below production threshold is taken into account as a **continuous energy loss**
- Similar approach is used for the **ionisation** process where spectrum of  $\chi$ -electrons is proportional to  $1/T^2$

Gamma Energy distribution (GeV)



22 27. Passage of particles through matter



# Cut and production thresholds for energy loss processes

- User defines cut in range expressed in units of length
- Using this range Geant4 kernel compute production threshold  $T_{cut}$  for each material during initialization
- For a typical process (G4hIonisation, G4eIonisation, ...), the production threshold  $T_{cut}$  subdivides the continuous and discrete parts of energy loss:

- Mean rate of energy lost due to soft energy transfers
- Total XS for discrete  $\gamma$ -electron production above  $T_{cut}$

$$\frac{dE(E, T_{cut})}{dx} = n_{at} \int_0^{T_{cut}} T \frac{d\sigma(Z, E, T)}{dT} dT$$

$$\sigma(Z, E, T_{cut}) = \int_{T_{cut}}^{T_{max}} \frac{d\sigma(Z, E, T)}{dT} dT$$


- At each step energy deposition is sampled by a fluctuation model using the computed mean energy loss
- Optionally, energy loss may be modified :
  - for the generation of extra  $\gamma$ -electrons under the threshold when the track is in the vicinity of a geometrical boundary (sub-cutoff)
  - for the sampling of fluorescence and Auger-electrons emission
- 4-momentum balance is provided in all cases



# Effect of production thresholds

500 MeV incident protons on EM Pb/LAr calorimeter

## In Geant4



One sets the production threshold for delta rays as a unique range:



1.5 mm

It is converted by Geant4 to energy:

$T_c = 455$  keV electron energy in liquid Ar

$T_c = 2$  MeV electron energy in Pb

## In Geant3



one has to set the cut for delta-rays (DCUTE) as an energy threshold

either to the Liquid Argon value, thus producing many small unnecessary d-rays in Pb,

or to the Pb value, thus killing the d-rays production everywhere

# What particles have cuts?

---

- Since Geant4 9.3 cuts are defined for
  - Gamma
  - Electron
  - Positron
  - Proton
- Cut for proton is used for all hadrons and ions by elastic scattering processes

# Which processes use cuts ?

---

- It is not mandatory to use cuts
- Energy thresholds for gamma are used in Bremsstrahlung
- Energy thresholds for electrons are used in ionisation and e+e- pair production processes
- Energy threshold for positrons is used in the e+e- pair production process
- Energy thresholds for gamma and electrons are used optionally (“ApplyCuts” options) in all discrete processes
  - Photoelectric effect, Compton, gamma conversion
- Energy threshold for protons are used in processes of elastic scattering for hadrons and ions defining the threshold for kinetic energy of nuclear recoil

# Comments

---

- Range cut approach was established for simulation of energy deposition inside solid or liquid media
  - Sampling and crystal calorimeters
  - Silicon tracking
- For specific user application, it may be revised, for example, by defining different cuts in range for electron and gamma
  - Gaseous detectors
  - Muon system
- Tracking cuts may be also used (saving some CPU) for simulation of penetration via shielding or for simulation in non-sensitive part of the apparatus
  - Astrophysics applications

# How to define cut?

---

- ◉ Using UI interface to geant4 kernel:
  - /run/setCut 0.1 mm
  - /run/setCutForAGivenParticle e- 10 um
- ◉ Implementing virtual method SetCuts() of G4VUserPhysicsList
- ◉ In Geant4 examples several different implementations of cut definition in user code are shown
  - Including user defined UI commands
  - \$G4INSTALL/examples/extended/electromagnetic

# Cuts per G4Region

---

- Uniform cut in range providing balanced simulation of particle transport in media with different density
- Requirements for precision in different part of complex geometry may be very different
  - Micron precision in tracking devices millimeter precision in calorimeters
  - Unique value of the cut in range may be not effective and not practical

# Cuts per G4Region

---

- Geometrical volumes may be assigned to G4Region
- By default the only one G4Region is created associated with the World volume
- If more than one G4Region is created it is possible to have different cut values

# How to define cut for G4Region?

---

- **Using UI interface to geant4 kernel:**
  - `/run/setCutForRegion VertexDetector 1 um`
- **Implementing virtual method `SetCuts()` of `G4VUserPhysicsList`**
- **Examples are available:**
  - `$G4INSTALL/examples/extended/electromagnetic`
  - `TestEm8` – simple gaseous detector
  - `TestEm9` – more complicate setup with tracker and muon detectors
- **To printout cut values and production thresholds use UI command:**
  - `/run/particle/dumpCutValues`



# Thank you for your attention!

---

